

Lasercooled radium monofluoride: A molecular all-in-one probe for new physics

T. A. Isaev* and R. Berger†

Clemens-Schöpf Institute, TU Darmstadt, Petersenstr. 22, 64287 Darmstadt, Germany

The particular advantages of using diatomic molecule radium monofluoride (RaF) as a versatile molecular probe for physics beyond the Standard Model are highlighted. i) RaF was previously suggested as being potentially amenable to direct cooling with lasers. As shown in the present work, RaF's energetically lowest electronically excited state is of $^2\Pi$ symmetry (in contrast to BaF), such that no low-lying $^2\Delta$ state prevents efficient optical cooling cycles. ii) The effective electric field acting on the unpaired electron in the electronic ground state of RaF is estimated larger than in YbF, from which the best restrictions on the electron electric dipole moment (eEDM) were obtained experimentally. iii) Favourable crossings of spin-rotational levels of opposite parity in external magnetic fields exist, which are important for the measurement of the nuclear anapole moment of nuclei with a valence neutron. Thus, RaF appears currently as one of the most attractive candidates for investigation of parity-odd as well as simultaneously parity- and time-reversal-odd interactions in the realms of molecular physics.

INTRODUCTION

The principle advantage of using heavy-atom, polar, diatomic molecules for searching for space parity violation interactions ($\mathcal{P}$ -odd interactions) and simultaneously space parity and time reversal violating forces ($\mathcal{P}$, $\mathcal{T}$ -odd forces) is known for more than 30 years. Nevertheless, only the latest generation of molecular experiments finally surpassed their atomic competitors in sensitivity to one of the most important $\mathcal{P}$, $\mathcal{T}$ -odd properties of elementary particles, namely the permanent electric dipole moment of an electron (eEDM) [1]. Whereas molecules are ideally tailored to create favourable, well-defined fields at the heavy nucleus, the complexity of measurements with molecules is typically connected with various systematic effects which can mimic $\mathcal{P}$ -odd correlations. This calls for an active identification of promising molecular candidates that allow different approaches to suppress systematic effects. In the search for an eEDM, experimentally oriented research groups are currently focussing on YbF [1], PbO [2], ThO [3], WC [4], PbF [5] molecules and HfH⁺ molecular ion [6]. In all these experiments, high-quality electronic structure calculations are crucial both for the preparation stage of the experiment and for subsequent interpretation of experimental data obtained [7].

Another set of pivotal molecular experiments is connected with attempts to measure the nuclear anapole moment, a $\mathcal{P}$ -odd electromagnetic form-factor appearing in $I > 0$ nuclei due to $\mathcal{P}$ -odd nuclear forces. The only nucleus, in which the anapole moment was successfully determined, is ^{133}Cs in experiment with a vapour of Cs atoms [8]. The result of this experiment is apparently in disagreement with the earlier measurement on Tl atom [9, 10]. Currently molecular experiments are under development in Yale on BaF [2] and in Groningen on SrF [11].

Recently, we identified the open-shell diatomic molecule RaF as an exceptionally suitable candidate for

nuclear anapole moment measurements, having on the one hand a high enhancement factor for nuclear spin-dependent weak interaction and on the other hand offering potential for direct cooling with lasers. In the present work we demonstrate that RaF presents unique possibilities for measurements of $\mathcal{P}$ -odd and $\mathcal{P}$, $\mathcal{T}$ -odd effects due to favourable combinations of peculiarities of molecular electronic structure and nuclear structure of isotopes of radium.

DIRECT COOLING OF MOLECULES WITH LASERS

We identified earlier a set of requirements on molecular electronic structure for molecules being suitable for direct cooling with lasers [12]. Monofluorides of group II elements (e.g. BaF and RaF) belong to the first class of molecules with highly diagonal Frank-Condon matrix. One problem emerging, however, in lasercooling of BaF molecule (isovalent to RaF) is connected with the existence of a metastable $^2\Delta$ level, lying energetically below the $^2\Pi$ level involved in the optical cooling loop. Our previous electron correlation calculations of the spectroscopic parameters in RaF indicated that the energetically lowest electronically excited level is $^2\Pi$. We took now larger atomic basis sets and active spaces of virtual molecular orbitals to investigate the stability of the ordering of electronic levels. The results are given in Table I. These show that even considerable alterations in the parameters of the Fock-space relativistic coupled cluster (FS-RCC as implemented in Dirac [13]) calculations do not change the ordering of levels in RaF. On the other hand FS-RCC, calculations of BaF with the basis set of similar quality as for RaF and as large active spaces (see supplementary material) confirm the first excited electronic level in BaF to be $^2\Delta$. Based on comparison to experimental BaF results we estimate the accuracy of $\tilde{T}_e$ calculation for RaF to be within 1200 cm^{-1} (without

changing the ordering of the levels), R_e within about $0.1 a_0$ and $\tilde{\omega}_e$ about 60 cm^{-1} .

LEVEL CROSSING IN MAGNETIC FIELD AND SENSITIVITY TO THE NUCLEAR ANAPOLE MOMENT

$\mathcal{P}$ -odd effects in diatomic molecules can be greatly enhanced by shifting levels of opposite parity to near-crossing with the help of external magnetic fields [17]. This idea is exploited in [2] for measurement attempts of the nuclear anapole moment in BaF. One of the main problems in the suggested approach is to create highly homogeneous magnetic fields in large volumes. Favourable values of magnetic fields required to tune spin-rotational levels to near crossing would be below 10 kG (1 T), as creation of larger magnetic flux densities require special effort, for instance superconducting magnets. To estimate if fields with $|B| < 1T$ suffice to create near level crossings in RaF we calculated Zeeman splittings for spin-rotational levels (See Fig. I and Fig. II in supplementary materials). Matrix elements of the spin-rotational Hamiltonian in magnetic field were implemented as in Ref. [18]. The following parameters of the spin-rotational Hamiltonian were employed: rotational constant $B_e = 5689 \text{ MHz}$ (as calculated from equilibrium structure), ratio of spin-doubling constant Δ to $2B$, $\Delta/2B = 0.97$ (as calculated within a four-component Dirac–Hartree–Fock approach), components of the hyperfine tensor for ^{225}Ra nucleus $A_{\parallel} = -15100 \text{ MHz}$ and $A_{\perp} = -14800 \text{ MHz}$ (calculated with the ZORA approach, values were already used for scaling in Ref. [19], but not explicitly reported therein) and components of G -tensor $G_{\parallel} = 1.993$ and $G_{\perp} = 1.961$ (crude estimate based on results for HgH). According to our calculation, the first crossing of levels of opposite parity takes place at about 3 kG for levels with the projection F_z of the total angular momentum on the direction of the magnetic field being $-3/2$. A few more crossings take place in fields up to 10 kG for levels with $F_z = -1/2$, thus providing additional freedom for choosing the optimal experimental parameters.

To estimate the lowest possible flux of RaF molecules, which allows to measure the anapole moment in RaF, one needs to find the ratio $\frac{\Delta W}{W}$ with W being the experimentally measured signal proportional to matrix element of the nuclear spin-dependent weak interaction W_a , and ΔW its experimental uncertainty. Of course the condition for meaningful measurement is $\frac{\Delta W}{W} < 1$. We assume that the experimental scheme employed for anapole moment measurement is analogous to the one suggested in [2]. According to [2] the maximal value of $\frac{\Delta W}{W}$ is

$$\frac{\Delta W}{W} \simeq \frac{1}{2\sqrt{2N_0}tW}, \quad (1)$$

with N_0 being the total number of molecules ($N_0 = F\tau$,

where F is *detected* molecular flux and τ is the total measurement time) and t the interaction time between molecule and external fields. Thus one gets $F > 1/(8W^2t^2\tau)$. The time for molecular trapping can reach a few seconds [20], so let $t = 1 \text{ s}$, $\tau = 1 \text{ h} = 3600 \text{ s}$. To estimate W we just scale the W value for Ba given in Ref. [2], as $W^{\text{Ba}}/W^{\text{Ra}} \simeq W_a^{\text{Ba}}/W_a^{\text{Ra}}$, and take $W^{\text{Ra}} \simeq 10W^{\text{Ba}} = 50 \text{ Hz}$. The least required flux of RaF is roughly then $F = 1/(8 \cdot 2500 \cdot 1 \cdot 3600) = 1.4 \cdot 10^{-8} \text{ s}^{-1}$. In practice one might expect trapping and detection of at least one molecule during an experiment time of $\tau = 1 \text{ h}$ (see below), which corresponds to the flux $2.8 \cdot 10^{-4} \text{ s}^{-1}$, which is a few orders of magnitude higher than the least required flux of RaF. In practice this implies that it should be possible to perform successful measurements with signals from *single* trapped RaF molecules.

EFFECTIVE ELECTRIC FIELD ACTING ON THE UNPAIRED ELECTRON IN RAF

One of the most important parameters in molecular experiments on eEDM is the effective electric field acting on the unpaired electron in the electronic ground state of the molecule of interest. This field, however, can not be measured directly in experiment, but is predicted from quantum chemical calculations. We estimate here the effective electric field acting on the unpaired electron in RaF by using relations between matrix elements of different $\mathcal{P}$ -odd and $\mathcal{P}, \mathcal{T}$ -odd operators as it has been done in Ref. [21] by Kozlov and extended recently in Ref. [22]. According to the semiempirical model of Kozlov [21] the relation between the parameter W_s of the $\mathcal{P}, \mathcal{T}$ -odd term and the parameter W_a of the $\mathcal{P}$ -odd term in the effective spin-rotational Hamiltonian is $W_s/W_a = Z3\gamma/(2\gamma + 1)$, where Z is the nuclear charge number and $\gamma = \sqrt{1 - (\alpha Z)^2}$.

On the other hand, matrix elements M_{SPT} of the scalar $\mathcal{P}, \mathcal{T}$ -odd interaction and matrix elements M_{EDM} of the coupling between eEDM and inner molecular electric field can be expressed as [22]:

$$\begin{aligned} M_{\text{SPT}}/C_{\text{SP}} &= \frac{G_F A}{3\sqrt{2}} \alpha A_s A_p \rho_0(A) R_N(A) (1 - 0.56Z^2\alpha^2/2) \\ M_{\text{EDM}}/d_e &= \frac{4Z^3\alpha^2}{\gamma(4\gamma^2 - 1)\nu_p^{3/2}\nu_s^{3/2}} \end{aligned} \quad (3)$$

In the equation above G_F is Fermi's constant, A the atomic mass number, $R_N(A) = 1.2A^{1/3} \text{ fm}$ the effective nuclear radius and $\rho_0(A) = 4\pi/(3R_N(A)^3)$. The parameters A_s and A_p are normalisation factors of the electronic wavefunction and can be expressed in the semiclassical approximation as [22]:

$$A_s = \frac{(1 + \gamma)(2ZR_N(A))^\gamma}{(1 - 0.375Z^2\alpha^2)\sqrt{Z}\Gamma(2\gamma + 1)\nu_s^{3/2}} \quad (4)$$

$$A_p = \frac{\sqrt{Z}(2ZR_N(A))^\gamma}{(1 - 0.375Z^2\alpha^2)\Gamma(2\gamma + 1)\nu_p^{3/2}}, \quad (5)$$

TABLE I. Estimated molecular spectroscopic parameters for the electronic ground state $^2\Sigma_{1/2}$ and the first electronically excited state from FS-CCSD calculations of ^{226}RaF and ^{137}BaF . 17 electrons are correlated in both cases: $2p_{1/2,3/2}$ and $2s_{1/2}$ on fluorine, $5s_{1/2}$, $5p_{1/2,3/2}$ and $6s_{1/2}$ and Ba and $6s_{1/2}$, $6p_{1/2,3/2}$ and $7s_{1/2}$ on Ra. The results are given for two different 4-component all-electron uncontracted basis sets on Ra and Ba atom: Dyall's relativistic basis set and the RCC-ANO basis set by Roos et al. The active space in FS-CCSD was restricted by energy for Dyall's basis set up to 10 Hartree and for RCC-ANO up to 1000 Hartree (see supplementary material for computational details and references).

	R_e/a_0	$\tilde{\omega}_e/\text{cm}^{-1}$	$\tilde{T}_e/\text{cm}^{-1}$	$\tilde{D}_e/\text{cm}^{-1}$
RaF				
Dyall basis set				
$^2\Sigma_{1/2}$	4.24 ^a	428 ^a		32104 ^a
$^2\Pi_{1/2}$	4.24 ^a	432 ^a	14012 ^a	31302 ^a
$^2\Pi_{3/2} + ^2\Delta_{3/2}$	4.25	410	15967	
$^2\Delta_{3/2} + ^2\Pi_{3/2}$	4.25	432	16435	
$^2\Delta_{5/2}$	4.27	419	17086	
$^2\Sigma_{1/2}$	4.26	416	18118	
RCC-ANO basis set				
$^2\Sigma_{1/2}$	4.29	431		52425
$^2\Pi_{1/2}$	4.29	428	13263	
$^2\Pi_{3/2} + ^2\Delta_{3/2}$	4.31	415	15020	
$^2\Delta_{3/2} + ^2\Pi_{3/2}$	4.28	431	15405	
$^2\Delta_{5/2}$	4.30	423	15840	
$^2\Sigma_{1/2}$	4.32	419	16682	
BaF				
RCC-ANO basis set				
$^2\Sigma_{1/2}$	4.15	456		45006
$^2\Delta_{3/2}$	4.21	455	10871	
$^2\Delta_{5/2}$	4.21	455	11304	
$^2\Pi_{1/2}$	4.19	446	11701	
$^2\Pi_{3/2}$	4.19	444	12354	
$^2\Sigma_{1/2}$	4.23	455	14220	
Experimental data				
(from Ref. [14] if not indicated otherwise)				
$^2\Sigma_{1/2}$	4.09 ^b	469 ^b		46780 ± 726^b
$^2\Delta$		437	10940	
$^2\Pi$	4.13 ^c	437	11727	
$^2\Sigma$	4.17 ^c	424	13828	

a) Ref. [12]; b) Ref. [15]; c) Ref. [16]

with $\Gamma(x)$ being the gamma-function and ν the effective principal quantum number. Upon relating these quantities, one has to take into account different definitions of the parameters of the effective Hamiltonian and $\mathcal{P}$, $\mathcal{T}$ -odd parameters (W_s and W_c) in [22] and [21] (see [22] for details).

It is interesting to note that in experiments with one kind of molecules the scalar $\mathcal{P}$, $\mathcal{T}$ -odd interaction is indistinguishable from the eEDM effect – they both contribute to $\mathcal{P}$, $\mathcal{T}$ -odd electron paramagnetic resonance signal. One can, however, separate the contributions by taking data from experiments with different molecules (or molecule and atom) like it is proposed in [22] for eEDM in Tl and YbF.

Taking into account the above relations one can easily estimate the effective electric field acting on the electron in the ground $^2\Sigma$ state of RaF. The results are given in Table II. The accuracy of such an estimate is not high, but sufficient to identify RaF as promising candidate for eEDM measurements.

Another attractive feature of measurements with Ra nuclei is that there exist also a nuclear mechanism enhancing $\mathcal{P}$ -odd and $\mathcal{P}$, $\mathcal{T}$ -odd effects in certain Ra isotopes. According to [25] and [26] the Schiff moment in nuclei possessing octapole deformation is enhanced by about 10 to 100 times. The mechanism is similar to the one in diatomic and chiral molecules: enhancement is reached due to closeness of rotational levels of opposite

TABLE II. Estimates of the $|W_s|$ parameter of $\mathcal{P}$, $\mathcal{T}$ -odd scalar interaction (in kHz) and effective field on the electron $|E_{\text{eff}}|$ (in units $10^{24} \text{ Hz e}^{-1} \text{ cm}^{-1}$) for open-shell diatomic molecules. For comparison the data of *ab initio* calculations of $|E_{\text{eff}}|$ are provided.

	Z	$ W_a /\text{Hz}^a)$	$ W_s /\text{kHz}$	$ E_{\text{eff}} $	$ E_{\text{eff}} $ (<i>ab initio</i>)
BaF	56	1.9×10^2	12	1.8	$1.9^b)$
YbF	70	6.1×10^2	38	4.1	$6.0^c)$
RaF	88	2.1×10^3	13×10^1	9.5	
HgH	80	2.0×10^3	19×10^1	16	
CnH	112	3.1×10^4	35×10^2	300	

a) Results of Ref. [19]

b) Results of Ref. [23]

c) Results of Ref. [24]

parity. As a result the estimated Schiff moments in ^{225}Ra and ^{223}Ra isotopes are equal to 300 and 400 (in units $\eta 10^8 \text{ e fm}^3$, where η is effective nucleon-nucleon $\mathcal{P}$, $\mathcal{T}$ -odd force constant), respectively, whereas for ^{199}Hg for example it is only -1.4 (data taken from Ref. [26]).

PRODUCTION OF RAF

Besides the possible routes discussed in Ref. [12], we propose in Ref. [27] to produce neutral RaF via RaF^+ , which is subsequently neutralised by charge exchange in collision with a suitably chosen collision gas or by interaction with surfaces that provide the adequate work function for an iso-enthalpic electron transfer. RaF^+ can in turn be formed in reactive collisions of radium ions with a suitable fluorine containing compound.

CONCLUSION

We demonstrated various special properties of RaF that render it a versatile molecular laboratory for studying wide range of physical phenomena, from laser cooling to the physics beyond the Standard Model. Unique combination of rovibronic, electronic and nuclear structure features render RaF particularly attractive for further experimental study. The authors are grateful to D. DeMille, V. Flambaum, M. Kozlov and S. Hoekstra for discussion.

* timur.isaev@tu-darmstadt.de

† robert.berger@tu-darmstadt.de

[1] J. J. Hudson, D. M. Kara, I. J. Smallman, B. E. Sauer, M. R. Tarbutt, and E. A. Hinds, *Nature* **473**, 493 (2011)

- [2] D. DeMille, S. B. Cahn, D. Murphree, D. A. Rahmlow, and M. G. Kozlov, *Phys. Rev. Lett.* **100**, 023003 (2008)
- [3] A. C. Vutha, B. Spaun, Y. V. Gurevich, N. R. Hutzler, E. Kirilov, J. M. Doyle, G. Gabrielse, and D. DeMille, *Phys. Rev. A* **84**, 034502 (2011)
- [4] J. Lee, E. R. Meyer, R. Paudel, J. L. Bohn, and A. E. Leanhardt, *J. Mod. Opt.* **56**, 2005 (2009)
- [5] N. E. Shafer-Ray, *Phys. Rev. A* **73**, 034102 (2006)
- [6] A. E. Leanhardt, J. L. Bohn, H. Loh, P. Maletinsky, E. R. Meyer, L. C. Sinclair, R. P. Stutz, and E. A. Cornell, *J. Mol. Spectrosc.* **270**, 125 (2011)
- [7] A. V. Titov, N. S. Mosyagin, A. N. Petrov, T. A. Isaev, and D. P. DeMille, *Progr. Theor. Chem. Phys.* **B 15**, 253 (2006)
- [8] C. S. Wood, S. C. Bennett, D. Cho, B. P. Masterson, J. L. Roberts, C. E. Tanner, and C. E. Wieman, *Science* **275**, 1759 (1997)
- [9] P. A. Vetter, D. M. Meekhof, P. K. Majumder, S. K. Lamoreaux, and E. N. Fortson, *Phys. Rev. Lett.* **74**, 2658 (1995)
- [10] J. S. M. Ginges and V. V. Flambaum, *Physics Reports-review Section of Physics Letters* **397**, 63 (2004)
- [11] J. E. van den Berg, S. H. Turkesteen, E. B. Prinsen, and S. Hoekstra, *Eur. Phys. J. D* **66** (2012)
- [12] T. A. Isaev, S. Hoekstra, and R. Berger, *Phys. Rev. A* **82**, 052521 (2010)
- [13] DIRAC, a relativistic ab initio electronic structure program, Release DIRAC08 (2008), written by L. Visscher, H. J. Aa. Jensen, and T. Saue, with new contributions from R. Bast, S. Dubillard, K. G. Dyall, U. Ekström, E. Eliav, T. Fleig, A. S. P. Gomes, T. U. Helgaker, J. Henriksson, M. Iliaš, Ch. R. Jacob, S. Knecht, P. Norman, J. Olsen, M. Pernpointner, K. Ruud, P. Salek, and J. Sikkema (see <http://dirac.chem.sdu.dk>)
- [14] C. Effantin, A. Bernard, J. Dincan, G. Wannous, J. Verges, and R. Barrow, *Molecular Physics* **70**, 735 (1990)
- [15] G. Blue, J. Margrave, T. Ehlert, and J. Green, *Nature* **199**, 804 (1963)
- [16] NIST database, “<http://webbook.nist.gov/chemistry/formser.html>,”
- [17] V. V. Flambaum and I. B. Khriplovich, *Phys. Lett. A* **110**, 121 (1985)
- [18] M. G. Kozlov, L. N. Labzowsky, and A. O. Mitrushchenkov, *Sov. Phys. JETP* **73**, 415 (1991)
- [19] T. A. Isaev and R. Berger, *Phys. Rev. A* **86**, 062515 (2012)
- [20] S. Hoekstra, J. J. Gilijamse, B. Sartakov, N. Vanhaecke, L. Scharfenberg, S. Y. T. van de Meerakker, and G. Meijer, *Phys. Rev. Lett.* **98** (2007)
- [21] M. G. Kozlov, *Sov. Phys. JETP* **62**, 1114 (1985)
- [22] V. A. Dzuba, V. V. Flambaum, and C. Harabati, *Phys. Rev. A* **84**, 052108 (2011)
- [23] M. G. Kozlov, A. V. Titov, N. S. Mosyagin, and P. V. Souchko, *Phys. Rev. A* **56**, R3326 (1997)
- [24] N. S. Mosyagin, M. G. Kozlov, and A. V. Titov, *J. Phys. B* **31**, L763 (1998)
- [25] N. Auerbach, V. V. Flambaum, and V. Spevak, *Phys. Rev. Lett.* **76**, 4316 (1996)
- [26] V. Spevak, N. Auerbach, and V. Flambaum, *Phys. Rev. C* **56**, 1357 (1997)
- [27] T. A. Isaev, S. Hoekstra, and R. Berger, “Ion neutralisation mass-spectrometry route to radium monofluoride (RaF),” to be published